

A-10-183-1

بررسی موتاسیون های ژن بتا تالاسمی در ۱۰۰ ناقل بیماری بتا تالاسمی در شمال شرق ایران

فرناز مهاجر تهران ۱*، کاظم قدسی ۲

۱- مشهد- دانشگاه علوم پزشکی مشهد- بیمارستان قائم - بخش ژنتیک, ۲- مشهد- دانشگاه علوم پزشکی مشهد- بیمارستان قائم - بخش ژنتیک,

mohajertf1@mums.ac.ir, kghodsi@yahoo.com

B-thalassemia gene mutations analysis in 100 carriers in North East of Iran 2006-2007
Farnaz Mohajer Tehran 1, Kazem Ghodsi 1, *MSc degree in Molecular Genetics, Position: Laboratory of Medical Genetics, Ghaem hospital, Mashhad University of Medical Sciences (MUMS), Iran Department of Medical Genetics, Ghaem Hospital, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran1. Introduction: Thalassemia is an inherited autosomal recessive blood disease. The -thalassemia can be caused by more than 200 HBB gene mutations on chromosome 11. Each population-at-risk, however, has its own spectrum of common mutations, usually from five to ten; a finding that simplifies mutation analysis and thus, determines the origin of the mutant genes. Commonly occurring mutations of the HBB gene are detected by a number of PCR-based procedures including reverse dot blot analysis or primer-specific amplification. Materials and Methods: Carriers of B-thalassemia gene mutation based on the hematological indices were referred to Molecular Genetics Laboratory in Ghaem Hospital. Standard questionnaire forms were filled out and pedigrees were drawn. So careful three-tier approach involving: (1) Full blood count (2) Special hematological tests, followed by (3) DNA mutation analysis using ARMS PCR, provides the most effective way in which to detect primary gene mutations. With the exception of deletions and rearrangements which were sequenced, the molecular lesions causing hemoglobinopathies are all identifiable by PCR-based techniques. Results: We have studied 200 chromosomes for mutations leading to -thalassemia. Gender study showed 48% patients were male and 52% were female. Average age was 25.7 years. In this study, IVSI-5 appeared to be the most common mutation with an occurrence rate of 33.3% among the 11 different mutations found to be associated with -thalassemia including CD44 13.7%, CD15 9.8%, IVS II-I 7.8%, IVS I-I 7.8%, CD8/9 7.8%, CD36/37 5.8%, CD30 3.9%, IVS1-6 1.9%, IVS1-110 1.9%, IVSII-654 1.9%. We have successfully detected 64% of the β -thalassemia mutations by using amplification refractory mutation system (ARMS) technique and 36% of the β -thalassemia mutations by using sequencing technique. Prenatal study showed that 31.5% were normal, 52.6% carrier, and 15.7% were affected. Pedigrees study demonstrated that 24% of patients had at least one B-thalassemia patients. Fifty six percent of marriages were first cousin. Hematological study revealed MCV 52- 77.5 fL, HgA2 2.3% -6.6% in carriers. Key word: Beta-thalassemia, carrier, Genetics screening, ARMS-PCR Email: mohajertf1@mums.ac.ir

واژه‌های کلیدی: بتا تالاسمی، ناقل، PCR-ARMS

B-thalassemia gene mutations analysis in 100 carriers in North East of Iran 2006-2007